arXiv:hep-ph/0009288v1 25 Sep 2000

FISIST/14-2000/CENTRA
September 2000

Particle Rapidity Density Saturation in Heavy Ion Collisions and the Dual String Model

J. Dias de Deus and R. Ugoccioni
CENTRA and Departamento de Física (I.S.T.),
Av. Rovisco Pais, 1049-001 Lisboa, Portugal

Abstract

We show that the Dual String Model with fusion leads, in heavy ion collisions, to strict saturation of the particle (pseudo-)rapidity density, normalised to the number of participant nucleons, as that number increases. Asymptotically, as $\sqrt{s} \to \infty$, with the number of participants fixed, this density approaches the nucleon-nucleon density.

A comparison with recent WA98 data is presented.

1 Introduction

Recently, experimental [1, 2, 3] and theoretical [4, 5] papers have addressed the question of the dependence of measurable quantities (charged particle density, transverse energy, J/ψ production rate) on the number $N_{\rm part}$ of participant nucleons in high energy heavy ion collisions. That information, as stressed in [4], as well as in [5], is extremely important as it allows for a better understanding of the initial conditions in the evolution of newly created dense matter and provides the information for discriminating among different models.

In the analysis of the data of [1] made in [4] it was emphasised the role of the dependence of the particle rapidity density, dN/dy, normalised to the number of pairs of participants, and two models were considered.

The HIJING Monte Carlo model [6] contains soft and semi-hard interactions in an unitarised form, the p_T threshold for jet production, p_0, being independent of energy and atomic mass number A. As in the model there is a growing number of hard collisions, growing faster than $N_{\rm part}$, the quantity

$$\Phi(N_{\rm part}, y, \sqrt{s}) \equiv \frac{1}{\frac{1}{2}N_{\rm part}}\frac{dN}{dy} \tag{1}$$

is an increasing function of $N_{\rm part}$, or

$$\frac{\partial\Phi}{\partial N_{\rm part}} > 0. \tag{2}$$

The EKRT model [7], on the other hand, relates the hard physics threshold p_s to a saturation criterion [8]: the number of partons multiplied by their effective area ($\approx 1/p_T^2$) must be less than the transverse area of interaction. The saturation transverse momentum p_s, in contrast to the parameter p_0 of the previous model, is a growing function of $\sqrt{s}$ and A. The assumption that physics is controlled by $p_T = p_s$ leads to the result that Φ is a *decreasing* function of $N_{\rm part}$,

$$\frac{\partial\Phi}{\partial N_{\rm part}} < 0. \tag{3}$$

In Fig. 1, taken from [4], we show the predictions for the two models, histogram for HIJING and dot-dashed line for EKRT.

The Dual String Model (DSM) with string fusion was developed in [5]. Concerning this model two remarks can be made:

(i) There exist two components, a valence-valence component, proportional to $N_{\rm part}$, and a sea-sea (including gluons) component which grows faster than $N_{\rm part}$ and corresponds to multiple scattering. We have a structure somewhat similar to the HIJING model. Without fusion DSM gives an increase of Φ as, roughly, $N_{\rm part}^{1/3}$.

(ii) With fusion, DSM leads to saturation of Φ as $N_{\rm part}$ increases, not very different from what is obtained in the EKRT model (see, in particular, [9]).

In conclusion, in DSM with fusion,

$$\frac{\partial\Phi}{\partial N_{\rm part}} > 0, \tag{4}$$



Figure 1: Central charged particle rapidity density per participating pair as a function of the number of participants. Results of HIJING (histograms), EKRT predictions (dot-dashed lines) and DSM predictions (solid lines) for central Au+Au collisions at $\sqrt{s} = 56, 130, 200$ AGeV. Also shown are results from pp and $p\bar{p}$ collisions and PHOBOS data (Everything in the figure except the DSM curves is taken from [4].)

as in HIJING, and, as N_{part} becomes larger and larger,

$$\frac{\partial \Phi}{\partial N_{\text{part}}} \xrightarrow[N_{\text{part}} \to \infty]{} 0, \tag{5}$$

The N_{part} dependence of Φ in DSM with fusion is also shown in Fig. 1 (solid lines). The parameters used are precisely those used in [5] to describe NA49 and PHOBOS data [1].

2 DSM and particle densities

DSM is essentially the Dual Parton Model [10] with the inclusion of strings [11]. The strings may interact by fusing [12] in the transverse plane of interaction and eventually one may reach a situation of percolation with the formation of extended regions of colour freedom, with the features of the expected Quark-Gluon Plasma.

In the Dual Parton Model hadrons are considered as made up of constituents quarks (valence and sea quarks) and gluons. Two basic diagrams contribute to particle production (see [5]). The valence-valence diagram corresponds to single inelastic scattering and the wounded nucleon model [13]. The sea-sea diagram (including gluons) corresponds to the additional inelastic multiple scattering contributions. These contributions may be internal, parton multiple scattering within the original valence-valence contribution, or external, involving other nucleons.

If h is the height of the valence-valence plateau, αh the height of the sea-sea plateau, $2k$ the average number of strings produced in a nucleon-nucleon collision we can write [5, 14] for the (pseudo-)rapidity particle density

$$\left.\frac{dN}{dy}\right|_{N_A N_A} = N_A\left[2+(2k-1)\alpha\right]h + (\nu_{N_A} - N_A)2k\alpha h, \tag{6}$$

where N_A is the number of pairs of participant nucleons

$$N_A = \frac{1}{2}N_{\text{part}}, \tag{7}$$

and ν_{N_A} is the average number of nucleon-nucleon collisions. From elementary multiple scattering arguments [14] we have

$$\nu_{N_A} = N_A^{4/3} \tag{8}$$

One should notice that the number of nucleon-nucleon collisions is $N_A + (\nu_{N_A} - N_A) = \nu_{N_A}$ and the number of strings is $N_A\left[2+2(k-1)\right] + (\nu_{N_A} - N_A)2k = 2k\nu_{N_A}$.

For $N_A = 1$, nucleon-nucleon collision, Eq. (6) gives

$$\left.\frac{dN}{dy}\right|_{pp} = \left[2+2(k-1)\alpha\right]h. \tag{9}$$

Assuming that h and α are energy independent (constant plateaus) the energy dependence of $dN/dy|_{pp}$ fixes the energy dependence of k.

From Eq.s (1), (6), (7), (8) and (9) we can write

$$\Phi(N_A, \sqrt{s}, y) = \phi(\sqrt{s}, y) + (N_A^{1/3} - 1)2k\alpha h \tag{10}$$

with

$$\phi(\sqrt{s}, y) \equiv \left.\frac{dN}{dy}\right|_{pp}. \tag{11}$$

The number of strings coming from nucleon multiple scattering —the second term in Eq. (6)—is $N_A(N_A^{1/3} - 1)2k$ and they occupy the transverse interaction area S_{N_A}, which, for central collisions, is approximately given by

$$S_{N_A} \simeq \pi\left(1.14 N_A^{1/3}\right)^2, \tag{12}$$

such that the dimensionless transverse density parameter η is

$$\eta = \left(\frac{r_s}{1.14}\right)^2 2k N_A^{1/3}(N_A^{1/3} - 1), \tag{13}$$

where $r_s \simeq 0.2$ fm is the string transverse section radius. Note that η increases with N_A and $\sqrt{s}$.

When fusion occurs, i.e., when strings cluster in the transverse plane, Eq. (10) becomes [5]

$$\Phi(N_A, \sqrt{s}, y) = \phi(\sqrt{s}, y) + F(\eta)(N_A^{1/3} - 1)2k\alpha h, \tag{14}$$

where $F(\eta)$ is the particle production reduction factor [15],

$$F(\eta) \simeq \sqrt{\frac{1 - e^{-\eta}}{\eta}}. \tag{15}$$

We shall next discuss the N_A, $\sqrt{s}$ and y dependence of Φ, Eq. (14).

For the N_A dependence we have

$$\frac{\partial \Phi}{\partial N_A^{1/3}} = 2k\alpha h \frac{\partial}{\partial N_A^{1/3}} \left[F(\eta)(N_A^{1/3} - 1) \right], \tag{16}$$

and, making use of the definition of η, Eq. (13),

$$\frac{\partial \Phi}{\partial N_A^{1/3}} = \frac{1.14}{r_s} \sqrt{2k} \alpha h \frac{\partial}{\partial N_A^{1/3}} \left[(1 - e^{-\eta}) \frac{N_A^{1/3} - 1}{N_A^{1/3}} \right]^{1/2}, \tag{17}$$

As both $(1 - e^{-\eta})$ and $(N_A^{1/3} - 1)/N_A^{1/3}$ are increasing functions of N_A, becoming constant as $N_A \to \infty$,

$$\frac{\partial \Phi}{\partial N_A} > 0 \qquad \text{with} \qquad \frac{\partial \Phi}{\partial N_A} \to 0 \quad \text{as} \quad N_A \to \infty. \tag{18}$$

These are the results announced in the Introduction, Eq.s (4) and (5).

The DSM with fusion thus predicts *saturation of the particle rapidity densities per participant pair of nucleons, Φ, as N_A increases.*

In Fig 2 we compare our model Eq. (14) with the very recent results [2] of WA98 Collaboration on central Pb+Pb collisions at the CERN SPS. For large values of N_{part} the saturation is very clearly seen, Fig. 2a, while for small values of N_{part} the rise of particle density with N_{part}, Fig. 2b, is well described. The values used for the parameters were $h = 0.77$ and $\alpha = 0.11$, and for the pp density we used the same value as WA98, from [16]. Note that the parameters here have slightly different values in comparison to the ones used in [5]. This is not surprising as the WA98 data are not compatible with NA49 data as given by PHOBOS [1].

Concerning the energy dependence of Eq. (14), if the energy is high enough

$$\phi(\sqrt{s}, y) \equiv \left. \frac{dN}{dy} \right|_{pp} \sim 2k\alpha h, \tag{19}$$

and

$$\Phi(N_A, \sqrt{s}, y)/\phi(\sqrt{s}, y) = 1 + F(\eta)(N_A^{1/3} - 1). \tag{20}$$



Figure 2: **(a)** Charged particle density per participant nucleon versus the number of participants; **(b)** absolute charged particle density versus the number of participants. The data from WA98 [2] refer to 158A GeV Pb+Pb collisions (filled circles), the open circle refers to pp collisions [16]; the solid line results from Eq. (14).

As $F(\eta)$ goes to zero as $\sqrt{s}$ increases we obtain the prediction that *asymptotically the particle density per participant, for a fixed number of participants, approaches the nucleon-nucleon density and this approach is controlled by the function* $F(\eta)$, Eq. (15). This means that asymptotically heavy ion collisions, with respect to particle densities, become similar to nucleon-nucleon collisions. This is a consequence of the increasing role played by string fusion.

In Fig. 3 we present our prediction for the dependence of the ratio (20) on the energy for fixed values of N_{part}.

Regarding the (pseudo-)rapidity dependence of Φ, Eq. (14), even without a more specific model, we can say that the valence-valence contribution dominates the large centre-of-mass $|y|$ region, as most of rapidity is taken by the valence diquarks. *In the fragmentation region we expect the particle density per participant nucleon to be equal to the nucleon-nucleon density,*

$$\Phi(N_A, \sqrt{s}, y_{\text{fragm}}) \simeq \phi(\sqrt{s}, y_{\text{fragm}}). \tag{21}$$

The sea-sea contributions are naturally shorter and more central in rapidity.

3 Conclusions

The DSM is a model with two components, the valence-valence component and the sea-sea component, the sea-sea component increasing its importance with energy and number



Figure 3: Ratio of the particle rapidity density per participating pair to the *pp* particle density, Eq. (20), as function of c.m. energy for fixed values of the number of participants, $N_{\rm part}$.

of participants. This is somewhat similar to the HIJING Monte Carlo model, with soft and hard components.

On the other hand, with fusion the DSM behaves, for large N_A, similarly to the EKRT model, but with strict saturation of the particle density per participant nucleon. However, in the original EKRT model the saturation criterion in the transverse plane is stronger than in case of fusion of strings. Here, saturation in the interaction area is asymptotic (when $\eta \to \infty$) while in the EKRT model it occurs, using the string language, when $\eta \equiv (\pi r_s^2)/(\pi 1.14^2 N_A^{2/3})(2kN_A(N_A^{1/3}-1)) = 1$. This causes the decrease of Φ with N_A in the EKRT original model.

Probably different explanations, such as the ones based on string fusion, parton saturation, parton shadowing, are in some sense dual and refer to the same underlying physics [14]. What is becoming clear is that saturation of particle density puts strong constraints in models, and limits the rise of the (pseudo-)rapidity plateau at RHIC and LHC.

Acknowledgements

R.U. gratefully acknowledges the financial support of the Fundação Ciência e Tecnologia via the “Sub-Programa Ciência e Tecnologia do 2^o Quadro Comunitário de Apoio.”

References

1. B.B. Back et al., PHOBOS Collaboration, preprint hep-ex/0007036, PHOBOS/RHIC.
2. M.M. Aggarwal et al., WA98 Collaboration, preprint nucl-ex/0008004, (WA98/CERN).

3. C. Lourenço, talk given at the Heavy Ion Day meeting in Lisbon, Portugal, April 2000.
4. X.-N. Wang and M. Gyulassy, preprint nucl-th/0008014.
5. J. Dias de Deus and R. Ugoccioni, preprint FISIST/11-2000/CENTRA (hep-ph/0008086), (I.S.T.), to appear in Phys. Lett. B.
6. X.-N. Wang and M. Gyulassy, Phys. Rev. D **44**, 3501 (1991); X.-N. Wang and M. Gyulassy, Comput. Phys. Commun. **83**, 307 (1994).
7. K.J. Eskola, K. Kajantie, P.V. Ruuskanen and K. Tuominen, Nucl. Phys. B **570**, 379 (2000).
8. L.V. Gribov, E.M. Levin and M.G. Ryskin, Physics Reports **100**, 1 (1983); A.H. Mueller and J. Qiu, Nucl. Phys. B **268**, 427 (1986).
9. K.J. Eskola, K. Kajantie and K. Tuominen, preprint JYFL-3700 and HIP-2000-45/TH (hep-ph/0009246), Univ. of Jyväskylä and Univ. of Helsinki.
10. A. Capella, U.P. Sukhatme, C.I. Tan and J. Trân Thanh Vân, Physics Reports **236**, 225 (1994).
11. N.S. Amelin, M.A. Braun and C. Pajares, Phys. Lett. B **306**, 312 (1993); N.S. Amelin, M.A. Braun and C. Pajares, Z. Phys. C **63**, 507 (1994).
12. N. Armesto, M.A. Braun, E.G. Ferreiro and C. Pajares, Phys. Rev. Lett. **77**, 3736 (1996); M. Nardi and H. Satz, Phys. Lett. B **442**, 14 (1998); J. Dias de Deus, R. Ugoccioni and A. Rodrigues, Eur. Phys. J. C **16**, 537 (2000).
13. A. Białas, B. Bleszyński and W. Czyż, Nucl. Phys. B **111**, 461 (1976).
14. N. Armesto and C. Pajares, Int. J. Mod. Phys. A **15**, 2019 (2000).
15. M.A. Braun and C. Pajares, Eur. Phys. J. C **16**, 349 (2000).
16. C. De Marzo et al., Phys. Rev. D **26**, 1019 (1982).